

**Open Access**

Meeting abstract

## **1005 Risk factors associated with increased coronary artery wall thickness by MRI: the multi-ethnic study of atherosclerosis (MESA)**

Robson Macedo<sup>\*1</sup>, Shaoguang Chen<sup>1</sup>, Shenghan Lai<sup>1</sup>, Steven Shea<sup>2</sup>, Kiang Liu<sup>3</sup>, Debiao Li<sup>3</sup>, Moyses Szklo<sup>4</sup>, Joao AC Lima<sup>1</sup> and David A Bluemke<sup>1</sup>

Address: <sup>1</sup>Johns Hopkins University|School Of Medicine, Baltimore, USA, <sup>2</sup>Siemens Medical Systems, Baltimore, USA, <sup>3</sup>Northwestern University, Chicago, IL, USA and <sup>4</sup>Johns Hopkins University School Of Public Health, Baltimore, USA

\* Corresponding author

from 11<sup>th</sup> Annual SCMR Scientific Sessions  
Los Angeles, CA, USA. 1–3 February 2008

Published: 22 October 2008

*Journal of Cardiovascular Magnetic Resonance* 2008, **10**(Suppl 1):A130 doi:10.1186/1532-429X-10-S1-A130

This abstract is available from: <http://jcmr-online.com/content/10/S1/A130>

© 2008 Macedo et al; licensee BioMed Central Ltd.

### **Introduction**

Coronary artery wall MRI has been shown to demonstrate increased thickness of the coronary wall in patients with clinically documented coronary artery disease.

### **Purpose**

The purpose of this study was to evaluate the relationship between coronary artery disease risk factors as well as imaging markers for atherosclerosis versus coronary artery wall thickness measured by MRI in asymptomatic individuals without known coronary artery disease.

### **Methods**

One hundred twenty three (123) asymptomatic subjects (57 males, mean age 62 +/- 9 yrs, 76 white, 46 blacks) in the Multi-Ethnic Study of Atherosclerosis (MESA) study were enrolled and underwent coronary wall MRI of the left and right coronary arteries at 5 mm intervals using a black blood turbo spin echo technique. All subjects had blood serum testing, blood pressure measurements, and anthropomorphic measurements. Carotid intimal-medial thickness and cardiac CT for calcium measurement were performed according to standard protocols.

### **Results**

In unadjusted models, male gender, diabetes, dyslipidemia, nonzero calcium score and common carotid intimal-medial thickness were associated with greater mean coronary artery wall thickness ( $p < 0.05$ ) (Table 1). In

multi-variable models of traditional risk factors (Model 1), male gender ( $p = 0.02$ ) remained significant and diabetes history showed a trend towards significance ( $p = 0.08$ ); coronary calcium score (zero vs. nonzero) or category of calcium score showed no significant relationship (model not shown). With the addition of common carotid IMT to Model 1, greater carotid IMT showed a strong relationship to coronary artery wall thickness (0.43 mm MRI/mm carotid IMT (0.11, 0.75),  $p = 0.009$ ) that was nearly unchanged with the addition of calcium score status (0.43 (0.11, 0.75),  $p = 0.008$ ) (Table 1, Model 2).

### **Conclusion**

In asymptomatic individuals free of clinical cardiovascular disease, greater coronary artery wall thickness by MRI was associated with gender, diabetes and dyslipidemia as well as computed tomography and ultrasound measures of atherosclerotic disease burden; overall coronary wall thickness was strongly related to common carotid IMT in multi-variable models.

**Table 1: In individuals free of clinical cardiovascular disease, coronary wall thickness by MRI was associated with gender, diabetes and dyslipidemia and computed tomography and ultrasound detected atherosclerotic disease; coronary wall thickness was strongly related to common carotid IMT in multi-variable models.**

Risk factor	Univariable Model		Multi-variable Model 1		Multi-variable Model 2	
	regression coefficient	p-value	regression coefficient	p-value	regression coefficient	p-value
Age (years)	0.0035 (-0.0021, 0.009)	0.22	0.0029 (-0.0028, 0.0087)	0.31	-0.0008 (-0.007, 0.0054)	0.8
Gender (ref, female)	0.14 (0.047, 0.24)	0.004	0.11 (0.017, 0.21)	0.02	0.070 (-0.028, 0.17)	0.16
Hypertension	0.042 (-0.056, 0.14)	0.4	-0.011 (-0.12, 0.095)	0.84	-0.036 (-0.14, 0.070)	0.5
Smoking history	0.032 (-0.065, 0.13)	0.51	0.028 (-0.069, 0.12)	0.57	0.03 (-0.065, 0.12)	0.53
Diabetes	0.19 (0.036, 0.3432)	0.02	0.14 (-0.015, 0.30)	0.08	0.097 (-0.059, 0.25)	0.22
Dyslipidemia	0.091 (-0.0045, 0.19)	0.06	0.054 (-0.048, 0.16)	0.29	0.061 (-0.040, 0.16)	0.24
CAC score > 0 (0 CAC = reference)	0.12 (0.023, 0.2123)	0.02			0.053 (-0.053, 0.16)	0.32
Carotid IMT (mm)	0.49 (0.21, 0.77)	0.0007			0.43 (0.11, 0.75)	0.009

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

